

Absorption of Spin of a Plane Circularly Polarized Wave

Radi I. Khrapko¹

Department of Physics, Moscow Aviation Institute, Moscow, Russia

Two current concepts of the electrodynamics spin are considered in comparison. Then the original concept, going back to Poynting and Sadowsky, was used to demonstrate the transfer of energy and spin from a plane circularly polarized electromagnetic wave into a symmetric absorber, i.e. into a medium, which is both dielectric and magnetic. A spin tensor is used. The given calculations show that spin is the same natural property of a plane electromagnetic wave, as energy and momentum. The expression for the torque density acting on the absorber is given.

Key Words: classical spin; circular polarization; spin tensor

PACS 75.10.Hk

1. Introduction 1. Spin density is proportional to the energy density

It was suggested as early as 1899 by Sadowsky [1] and as 1909 by Poynting [2], that any usual circularly polarized light carries angular momentum volume *density*, and the angular momentum density is proportional to the energy volume density. That is the angular momentum is present in any point of the light.

J.H. Poynting: If we put E for the energy in unit volume and G for the torque per unit area, we have $G = E\lambda / 2\pi$ [2, p. 565].

This sentence points that any absorption of a circularly polarized light results in a mechanical torque density acting on the absorber. We have researched this effect and have found that this torque density induces specific mechanical stresses in the absorber [3].

According to the Lagrange formalism, this angular momentum volume density is *spin density*. The spin of electromagnetic waves is described now by a canonical spin tensor [4 – 7].

$$Y_c^{\lambda\mu\nu} = -2A^{[\lambda} \delta_\alpha^{\mu]} \frac{\partial L}{\partial(\partial_\nu A_\alpha)}, \quad (1.1)$$

where $L = -F_{\mu\nu} F^{\mu\nu} / 4$ is the canonical Lagrangian, A^λ is the magnetic vector potential of the electromagnetic field, and $F_{\mu\nu}$ is the electromagnetic field tensor. So, any infinitesimal 3-volume dV_ν contains spin

$$dS^{\lambda\mu} = Y_c^{\lambda\mu\nu} dV_\nu. \quad (1.2)$$

The spin tensor (1.1) is appeared in the company of the energy-momentum tensor $T^{\mu\nu}$ and of the orbital angular momentum tensor, which is simply a moment of the energy-momentum tensor $2x^{[\lambda} T^{\mu]\nu}$. So, the total angular momentum tensor $J^{\lambda\mu\nu}$ equals the sum of orbital and spin angular momentums. The energy-momentum, total angular momentum, and spin tensors are

$$T_c^{\mu\nu} = -\partial^\mu A_\alpha F^{\nu\alpha} + g^{\mu\nu} F_{\alpha\beta} F^{\alpha\beta} / 4, \quad (1.3)$$

$$J_c^{\lambda\mu\nu} = 2x^{[\lambda} T_c^{\mu]\nu} + Y_c^{\lambda\mu\nu} = 2x^{[\lambda} T_c^{\mu]\nu} - 2A^{[\lambda} F^{\mu]\nu}, \quad Y_c^{\lambda\mu\nu} = -2A^{[\lambda} F^{\mu]\nu}. \quad (1.4)$$

So, the classical electromagnetic field theory provides meaningful descriptions of the spin and the orbital angular momentum separately. Note, spin tensor $Y^{\lambda\mu\nu}$ is not a moment of energy-

¹ Email address: khrapko_ri@hotmail.com, khrapko_ri@mai.ru <http://khrapori.wmsite.ru>

momentum tensor $2x^{[\lambda}T^{\mu]\nu}$, and spin density \mathbf{Y} is not (a part of) a moment of linear momentum density $\mathbf{r} \times (\mathbf{E} \times \mathbf{H})/c^2$.

According to the definition (1.4), the spin volume density is given by the component of the spin tensor

$$Y_c^{ij} = -2A^{[i}F^{j]l} = -2A^{[i}D^{j]l} = \mathbf{D} \times \mathbf{A} \quad (1.5)$$

For example, **Soper** [5] writes:

“To describe a circularly polarized plane wave traveling in the z-direction, we can choose a potential

$$A^x = a \cos[\omega(z-t)], A^y = -a \sin[\omega(z-t)] \quad (9.3.17).$$

The corresponding electric field is $E^k = -\partial_t A^k$.

$$E^x = -a\omega \sin[\omega(z-t)], E^y = -a\omega \cos[\omega(z-t)] \quad (9.3.18).$$

Thus the spin density carried by this wave is

$$\mathbf{s} = a^2 \omega \hat{\mathbf{z}} \quad (9.3.19),$$

where $\hat{\mathbf{z}}$ is a unit vector pointing in the z-direction.”

Soper means

$$Y_c^{xy} = E^x A^y - E^y A^x = a^2 \omega. \quad (1.6)$$

in agreement with (1.6).

Note that perfect plane electromagnetic waves were used by **Einstein** [8, § 7]:

“In the system K, very far from the origin of co-ordinates, let there be a source of electrodynamic waves, which in a part of space containing the origin of co-ordinates may be represented to a sufficient degree of approximation by the equations” of type (1.5).

The canonical spin tensor (1.4) was successfully used in [9] in order to confirm the fulfillment of the conservation laws with respect to spin when a plane circularly polarized electromagnetic wave reflects from a moving mirror, and in [10] where the absorption of spin of a plane wave is described. These calculations prove the functionality of the spin tensor and show that spin is the same natural property of a perfect plane electromagnetic wave, as energy and momentum; and spin density is proportional to energy density.

The classical experiments [11 – 14] also confirm that the spin density of plane waves is proportional to energy density. In these experiments, the angular momentum of the light was transferred to a half-wave plate, which rotated. So, work was performed *in any point of the plate*. This (positive or negative) amount of work reappeared as an alteration in the energy of the photons, i.e., in the frequency of the light, which resulted in moving fringes *in any point* of the interference pattern in a suitable interference experiment.

Some textbooks and articles point that infinite plane circularly polarized electromagnetic wave carries angular momentum:

F.S. Crawford, Jr.: “A circularly polarized travelling plane wave carries angular momentum” [15, p. 365].

R. Feynman: “... the photons of light that are right circularly polarized carry an angular momentum of one unit along the z-axis ... light which is right circularly polarized carries an energy and angular momentum” [16].

K. Bliokh and F. Nori: “... the plane wave carries the spin AM density \mathbf{S} defined as the *local* expectation value of the operator $\hat{\mathbf{S}}$ ” [17, p.4].

A spin tensor is used when describing plane waves in [18-23] and in other papers of the author.

2. Introduction 2. Is the spin density proportional to gradient of the energy density?

However, since 1939, another concept of electrodynamic spin is in use. The point is, Belinfante & Rosenfeld added specific terms,

$$\partial_\alpha (A^\mu F^{\nu\alpha}) \text{ and } 2\partial_\alpha (x^{[\lambda} A^{\mu]} F^{\nu\alpha}), \quad (2.1)$$

to the canonical energy-momentum tensor $T_c^{\mu\nu}$ and to the total angular momentum tensor $J_c^{\lambda\mu\nu}$, respectively [24,25], [5,Sec.9.4]. This procedure yields an energy-momentum tensor $T_{st}^{\mu\nu}$ (which differs from the Maxwell tensor $T^{\mu\nu}$), and an total angular momentum tensor $J_{st}^{\lambda\mu\nu}$. We named these tensors (2.2) and (2.3) "standard" [3]:

$$\begin{aligned} T_{st}^{\mu\nu} &= T_c^{\mu\nu} + \partial_\alpha (A^\mu F^{\nu\alpha}) = -\partial^\mu A_\alpha F^{\nu\alpha} + g^{\mu\nu} F_{\alpha\beta} F^{\alpha\beta} / 4 + \partial_\alpha (A^\mu F^{\nu\alpha}) \\ &= -\partial^\mu A_\alpha F^{\nu\alpha} + g^{\mu\nu} F_{\alpha\beta} F^{\alpha\beta} / 4 + \partial_\alpha A^\mu F^{\nu\alpha} + A^\mu \partial_\alpha F^{\nu\alpha} \\ &= g^{\mu\lambda} (-\partial_\lambda A_\alpha F^{\nu\alpha} + \partial_\alpha A_\lambda F^{\nu\alpha}) + g^{\mu\nu} F_{\alpha\beta} F^{\alpha\beta} / 4 + A^\mu \partial_\alpha F^{\nu\alpha} \\ &= -g^{\mu\lambda} F_{\lambda\alpha} F^{\nu\alpha} + g^{\mu\nu} F_{\alpha\beta} F^{\alpha\beta} / 4 + A^\mu \partial_\alpha F^{\nu\alpha} = T^{\mu\nu} + A^\mu \partial_\alpha F^{\nu\alpha}, \end{aligned} \quad (2.2)$$

where $T^{\mu\nu} = -g^{\mu\lambda} F_{\lambda\alpha} F^{\nu\alpha} + g^{\mu\nu} F_{\alpha\beta} F^{\alpha\beta} / 4$ is the Maxwell tensor,

$$\begin{aligned} J_{st}^{\lambda\mu\nu} &= J_c^{\lambda\mu\nu} + 2\partial_\alpha (x^{[\lambda} A^{\mu]} F^{\nu\alpha}) = 2x^{[\lambda} T_c^{\mu]\nu} - 2A^{[\lambda} F^{\mu]\nu} + 2\partial_\alpha (x^{[\lambda} A^{\mu]} F^{\nu\alpha}) \\ &= 2x^{[\lambda} T_c^{\mu]\nu} - 2A^{[\lambda} F^{\mu]\nu} + 2\delta_\alpha^{[\lambda} A^{\mu]} F^{\nu\alpha} + 2x^{[\lambda} \partial_\alpha (A^{\mu]} F^{\nu\alpha}) = 2x^{[\lambda} T_{st}^{\mu]\nu}. \end{aligned} \quad (2.3)$$

But this procedure eliminates spin tensor ($Y_{st}^{\lambda\mu\nu} = 0$). Really, the standard total angular momentum tensor (2.3) is equal to moment of the standard energy-momentum tensor: $J_{st}^{\lambda\mu\nu} = 2x^{[\lambda} T_{st}^{\mu]\nu}$ without a spin term, cf. (1.4)

So, the corresponding ("standard") spin term is absent. As a result, in the absence of electrodynamics spin tensor, it has been declared that the electrodynamics spin \mathbf{S} is contained inside the moment of linear momentum together with the orbital momentum \mathbf{L} [26, p. 7]

$$\int dV \epsilon_0 \mathbf{r} \times (\mathbf{E} \times \mathbf{B}) = \mathbf{L} + \mathbf{S},$$

and that a plane wave has no angular momentum at all.

Heitler W: "A plane wave travelling in z-direction and with infinite extension in the xy-directions can have no angular momentum about the z-axis, because $(\mathbf{E} \times \mathbf{B})$ is in the z-direction and $(\mathbf{r} \times (\mathbf{E} \times \mathbf{B}))_z = 0$ " [27].

According to the nowadays conception, electrodynamics spin density is proportional to *gradient of energy density*, not to the energy density itself:

Allen L., Padgett M. J.: "... the local spin angular momentum density per photon is proportional to the radial intensity gradient of a light beam:

$$j_z = -\frac{r}{2u^2} \frac{\partial(u^2)}{\partial r} \hbar\sigma$$

where $\sigma = \pm 1$ for right- and left-handed circularly polarized light respectively, u^2 is the beam intensity, and r is the distance from the axis. For a plane wave there is no gradient and the spin density is zero." [28]

Simmonds J. W., Guttman M. J.: "The electric and magnetic fields can have a nonzero z-component only within the skin region of this wave. Having z-components within this region implies the possibility of a nonzero z-component of angular momentum within this region.

So, the skin region is the only in which the z-component of angular momentum does not vanish" [29, p. 227]

Thus, according to the widespread opinion, an electromagnetic wave without an azimuthal phase gradient has an angular momentum only where, there is an intensity gradient, and this angular momentum is spin. Particularly, angular momentum of a circularly polarized beam is not an orbital angular momentum, but is a spin, and this spin is localized on the surface of the beam. This opinion carries the spin far to the edge of a real wave in compliance with the Humblet transformation [30].

We have criticized [3] the Humblet transformation. We have noted [31] that this concept, "Spin is only in the skin region," threatens us with a considerable nonlocality of the

electrodynamics because the concept implies that the energy and momentum of photons are absorbed everywhere in the absorber, but the spin is absorbed in the remote boundary of the wave.

Note, the Belinfante-Rosenfeld procedure does not yield the Maxwell tensor and even does not symmetrize the canonical tensor (1.3); the term $A^\mu \partial_\alpha F^{\nu\alpha}$ in (2.2) vanishes only when interactions are absent, but this case has no sense.

In this paper, a new calculation confirms the Poynting's and Sadowsky's concept. Absorption of spin by a conducting medium was considered in [23,32]. Now we consider a plane circularly polarized electromagnetic wave, which is absorbed in a "symmetric absorber".

3. A symmetric absorber

We call "symmetric absorber" a medium, which is both dielectric and magnetic with $\varepsilon = \mu$. Such a medium does not require generating a reflected wave; this simplifies formulas.

So, let a perfect plane monochromatic circularly polarized electromagnetic wave

$$F_{\alpha\beta} = \{E_x = 1, E_y = i, B^x = -i/c, B^y = 1/c\} \exp(ikz - i\omega t) E, \quad ck = \omega \quad (3.1)$$

$$F^{\mu\nu} = \{D^x = \varepsilon_0, D^y = i\varepsilon_0, H_x = -i\sqrt{\varepsilon_0/\mu_0}, H_y = \sqrt{\varepsilon_0/\mu_0}\} \exp(ikz - i\omega t) E, \quad (3.2)$$

impinges normally on a flat x,y-surface of the absorber, which is characterized by complex permittivity and permeability $\tilde{\varepsilon} = \tilde{\mu}$ (we indicate complex numbers by the *breve* mark).

So the wave propagated in the absorber is described by the formulas

$$F_{\alpha\beta} = \{E_x = 1, E_y = i, B^x = -i\tilde{\varepsilon}/c, B^y = \tilde{\varepsilon}/c\} \exp(i\tilde{k}z - i\omega t) E, \\ c\tilde{k} = \tilde{\varepsilon}\omega, \quad \tilde{k} = k_1 + ik_2, \quad \tilde{\varepsilon} = \varepsilon_1 + i\varepsilon_2 \quad (3.3)$$

$$F^{\mu\nu} = \{D^x = \tilde{\varepsilon}\varepsilon_0, D^y = i\tilde{\varepsilon}\varepsilon_0, H_x = -i\sqrt{\varepsilon_0/\mu_0}, H_y = \sqrt{\varepsilon_0/\mu_0}\} \exp(i\tilde{k}z - i\omega t) E, \quad (3.4)$$

4. Energy and spin flux

Using the Maxwell tensor $T^{\mu\nu} = -g^{\mu\lambda} F_{\lambda\alpha} F^{\nu\alpha} + g^{\mu\nu} F_{\alpha\beta} F^{\alpha\beta} / 4$ yields the Poynting vector in the vacuum

$$\langle c^2 T^{iz} \rangle = \Re\{-\bar{F}_{ix} F^{zx} + \bar{F}_{iy} F^{zy}\} / 2 = \mathbf{E} \times \mathbf{H} = \sqrt{\varepsilon_0/\mu_0} E^2 \quad [\text{J/m}^2\text{s}] \quad (4.1)$$

and the Poynting vector in the absorber

$$\mathbf{E} \times \mathbf{H} = \sqrt{\varepsilon_0/\mu_0} \exp(-2k_2 z) E^2. \quad (4.2)$$

Power volume density of the released energy in the absorber is

$$w = -\partial_z (\mathbf{E} \times \mathbf{H}) = 2k_2 \sqrt{\varepsilon_0/\mu_0} \exp(-2k_2 z) E^2 \quad [\text{J/m}^3\text{s}]. \quad (4.3)$$

Using the canonical spin tensor (1.4) $Y_c^{\lambda\mu\nu} = -2A^{[\lambda} F^{\mu]\nu}$ and $A_i = -\int E_i dt = -iE_i / \omega$ yields the spin flux density in the vacuum

$$\langle Y_c^{xyz} \rangle = \Re\{-\bar{A}^x F^{yz} + \bar{A}^y F^{xz}\} / 2 = \Re\{\bar{A}_x H_x + \bar{A}_y H_y\} / 2 = \sqrt{\varepsilon_0/\mu_0} E^2 / \omega \quad [\text{Js/m}^2\text{s}]. \quad (4.4)$$

and the spin flux density in the absorber

$$\langle Y_c^{xyz} \rangle = \Re\{-\bar{A}^x F^{yz} + \bar{A}^y F^{xz}\} / 2 = \Re\{\bar{A}_x H_x + \bar{A}_y H_y\} / 2 = \sqrt{\varepsilon_0/\mu_0} \exp(-2k_2 z) E^2 / \omega. \quad (4.5)$$

Torque volume density from the absorbed spin angular momentum in the absorber is

$$\tau_z = -\partial_z \langle Y_c^{xyz} \rangle = 2k_2 \sqrt{\varepsilon_0/\mu_0} \exp(-2k_2 z) E^2 / \omega \quad [\text{J/m}^3]. \quad (4.6)$$

4. Conclusion

The given calculations show that spin is a natural property of a plane electromagnetic wave, similar to energy. The Poynting's statement is confirmed: the absorption of energy of a circularly polarized light is accompanied by the volume density of torque.

We are eternally grateful to Professor Robert Romer, having courageously published the question: "Does a plane wave really not carry spin?" [33].

References

1. Sadowsky A. *Acta et Comm. Imp. Universitatis Jurievensis* 7, No. 1-3 (1899)
2. Poynting J. H., "The wave motion of a revolving shaft, and a suggestion as to the angular momentum in a beam of circularly polarised light". *Proc. R. Soc. Lond. A* 82, 560-567 (1909)
3. Khrapko R.I. "Mechanical stresses produced by a light beam" *J. Modern Optics*, **55**, 1487-1500 (2008) <http://khrapkori.wmsite.ru/ftpgetfile.php?id=9&module=files>
4. Corson E M *Introduction to tensors, spinors, and reativistic wave-equation* NY, Hafner, 1953 p.71
5. Soper D. E., *Classical Field Theory* (N.Y.: Dover, 2008), p. 114
6. Barut A. O. *Electrodynamics and Classical Theory of Particles and Fields* (Macmillan, New York, 1964), p. 102
7. Goedecke G. H. On stress-energy tensors *J. Math. Phys.* **15**, 792 (1974)
8. Einstein A. Zur Elektrodynamik bewegter Korper *Annalen der Physik* **17**, 891, 1905
9. Khrapko R.I. "Reflection of light from a moving mirror" *Optik* **136** (2017) 503–506 <http://khrapkori.wmsite.ru/ftpgetfile.php?id=153&module=files>
10. Khrapko R.I. "Absorption of angular momentum of a plane wave" *Optik* **154** (2018) 806-810 <http://khrapkori.wmsite.ru/ftpgetfile.php?id=161&module=files>
11. Righi A., *Mem. d. Accad. d. Scienze di Bologna*, 4 (1882), 247
12. Atkinson R., *Phys. Rev.*, 47 (1935), 623
13. Henriot E, *Comptes rendus* 198, 1146 (1934).
14. Wood R., *Physical Optics*, 2nd ed. (1911), p. 341.
15. Crawford F.S., Jr., *Waves: Berkley Physics Course - V. 3* (Berkeley, California June, 1968)
16. Feynman R. P., R. B. Leighton, M. Sands, *The Feynman Lectures on Physics* (Addison–Wesley, London, 1965) Vol. 3, p. 17–10.
17. K. Y. Bliokh, F. Nori, "Transverse and longitudinal angular momenta of light", *Physics Reports*, 592, 1-38 (2015).
18. Khrapko R.I. True energy-momentum tensors are unique. Electrodynamics spin tensor is not zero [arXiv:physics/0102084](https://arxiv.org/abs/physics/0102084)
19. Khrapko R.I. Violation of the gauge equivalence [arXiv:physics/0105031](https://arxiv.org/abs/physics/0105031)
20. Khrapko R.I. Classical spin in space with and without torsion" *Gravitation & Cosmology*, 10, No. 1-2, 91-98 (2004) <http://khrapkori.wmsite.ru/ftpgetfile.php?id=18&module=files>
21. Spin of the Classical Electrodynamics *Bulletin of Peoples' Friendship University of Russia. Series Physic.* 2002, #10(1), p. 40
22. Energy-momentum Localization and Spin". *Bulletin of Peoples' Friendship University of Russia. Series Physic.* 2002, #10(1), p. 35 <http://khrapkori.wmsite.ru/ftpgetfile.php?module=files&id=32>
23. Absorption of Spin by a Conducting Medium *AASCIT Journal of Physics* Vol. 4, No. 2, Page: 59-63 (2018) <http://www.aascit.org/journal/archive2?journalId=977&paperId=6355>
24. Belinfante F. J., *Physica* **6**, 887 (1939).
25. Rosenfeld L., "Sur le Tenseur D'Impulsion-Energie." *Memoires de l'Academie Royale des Sciences de Belgiques* **8** No 6 (1940).
26. Andrews D.L., M. Babiker (Editors) *The angular momentum of light* (Cambrige 2013)
27. Heitler W *The Quantum Theory of Radiation* (Oxford: Clarendon, 1954) p. 401.
28. Allen L., Padgett M. J. "Response to Question #79. Does a plane wave carry spin angular momentum?" *Am. J. Phys.* **70**, 567 (2002).
29. Simmonds J. W., M. J. Guttman, *States, Waves and Photons* (Addison-Wesley, Reading, MA, 1970)
30. Humblet J., "Sur le moment d'impulsion d'une onde electromagnetique". *Physica (Utrecht)* **10** (7): 585 (1943)

31. Khrapko R.I. "On the possibility of an experiment on 'nonlocality' of electrodynamics", *Quantum Electron.* 2012, **42** (12), 1133–1136.
<http://khrapkori.wmsite.ru/ftpgetfile.php?id=34&module=files>
32. Khrapko R I. "Absorption of light by a conducting medium" (2004)
<http://mai.ru/science/trudy/published.php?ID=34247> In Russian.
33. Khrapko R I "Does plane wave not carry a spin?" *Amer. J. Phys.* 69, 405 (2001)

Поглощение спина плоской волны

Р. И. Храпко

Московский авиационный институт, Москва, 125993

В сравнении рассмотрены две современные концепции спина электродинамики. Затем первоначальная концепция, восходящая к Пойнтингу и Садовскому, использовалась для демонстрации переноса энергии и спина из плоской электромагнитной волны круговой поляризации поляризованной в «симметричный поглотитель» (являющийся одновременно диэлектриком и магнетиком). Использован тензор спина. Эти расчеты показывают, что спин является таким же естественным свойством плоской электромагнитной волны как энергии и импульса. Приводится выражение для плотности момента силы, действующего на поглотитель.

Классический спин, круговая поляризация, тензор спина